

Low urine output

Normal urine output

- typically at least 0.5 cc/kg/hr.
- Oliguria: urine output < 400cc/day
- Anuria : urine output < 100cc/day

Do you believe the numbers?

- If patient has foley, flush tubing to make sure it is not clogged.
- If no foley, ask about urine output, look at daily weights, etc.

Examine patient bedside.

- Assess volume status: mucous membranes, skin pallor/dryness, edema, complaints of thirst, neck veins (assess CVP), crackles in lungs (pulmonary edema), bladder palpable on exam, prostate exam, etc.

Check a post-void residual by bladder scan.

- If volume > 300cc, then insert foley (In and out). If consistent > 300cc, keep foley in.

Causes of urinary retention:

- BPH, anticholinergic medication, side effect of medication: narcotics/benadryl/anesthetics

Assess for renal failure (AKI)

- Prerenal, renal, postrenal causes.
- Look for fluid overload (CHF), and obstruction (renal US).
- If both negative, then fluid challenge is acceptable: 500cc L bolus. (Go to AKI for further management).

CHF/Volume overloaded? Initiate diuresis:

- Working kidneys: lower IVF rate and self-diuresis
 - CHF/symptomatic: use lasix 20mg - 80mg IV
 - Renal failure: Dialysis? If kidneys still working, can try high dose lasix - 160mg- 240mg IV Lasix.
-

Revision #4

Created 25 February 2022 06:12:31 by Katarina Soewono

Updated 3 June 2022 02:48:54 by Katarina Soewono