

# Peds Immunization

Can also download cdc vaccination schedule app via your phone store

**Table 1** Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2022

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2).

Vaccine	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19–23 mos	2–3 yrs	4–6 yrs	7–10 yrs	11–12 yrs	13–15 yrs	16 yrs	17–18 yrs
Hepatitis B (HepB)	1 <sup>st</sup> dose	← 2 <sup>nd</sup> dose →			← 3 <sup>rd</sup> dose →												
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	See Notes												
Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose			← 4 <sup>th</sup> dose →				5 <sup>th</sup> dose					
Haemophilus influenzae type b (Hib)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	See Notes		← 3 <sup>rd</sup> or 4 <sup>th</sup> dose, See Notes →										
Pneumococcal conjugate (PCV13)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose		← 4 <sup>th</sup> dose →										
Inactivated poliovirus (IPV <18 yrs)			1 <sup>st</sup> dose	2 <sup>nd</sup> dose			← 3 <sup>rd</sup> dose →					4 <sup>th</sup> dose					
Influenza (IIV4) or Influenza (LAIV4)												Annual vaccination 1 or 2 doses					Annual vaccination 1 dose only
Measles, mumps, rubella (MMR)					See Notes		← 1 <sup>st</sup> dose →					2 <sup>nd</sup> dose					
Varicella (VAR)							← 1 <sup>st</sup> dose →					2 <sup>nd</sup> dose					
Hepatitis A (HepA)					See Notes												
Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)															1 dose		
Human papillomavirus (HPV)															See Notes		
Meningococcal (MenACWY-D ≥9 mos, MenACWY-CRM ≥2 mos, MenACWY-TT ≥2 years)															1 <sup>st</sup> dose	2 <sup>nd</sup> dose	
Meningococcal B (MenB-4C, MenB-FHbp)																	
Pneumococcal polysaccharide (PPSV23)																	
Dengue (DEN4CYD; 9–16 yrs)																	Seropositive in endemic areas only (See Notes)

Range of recommended ages for all children
  Range of recommended ages for catch-up vaccination
  Range of recommended ages for certain high-risk groups
  Recommended vaccination can begin in this age group
  Recommended vaccination based on shared clinical decision-making
  No recommendation/not applicable

## Table 2 Recommended Catch-up Immunization Schedule for Children and Adolescents Who Start Late or Who Are More than 1 Month Behind, United States, 2022

The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age. **Always use this table in conjunction with Table 1 and the Notes that follow.**

Children age 4 months through 6 years					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B	Birth	<b>4 weeks</b>	<b>8 weeks and at least 16 weeks after first dose</b> minimum age for the final dose is 24 weeks		
Rotavirus	6 weeks Maximum age for first dose is 14 weeks, 6 days.	<b>4 weeks</b>	<b>4 weeks</b> maximum age for final dose is 8 months, 0 days		
Diphtheria, tetanus, and acellular pertussis	6 weeks	<b>4 weeks</b>	<b>4 weeks</b>	<b>6 months</b>	<b>6 months</b>
Haemophilus influenzae type b	6 weeks	<b>No further doses needed</b> if first dose was administered at age 15 months or older. <b>4 weeks</b> if first dose was administered before the 1 <sup>st</sup> birthday. <b>8 weeks (as final dose)</b> if first dose was administered at age 12 through 14 months.	<b>No further doses needed</b> if previous dose was administered at age 15 months or older <b>4 weeks</b> if current age is younger than 12 months <b>and</b> first dose was administered at younger than age 7 months <b>and</b> at least 1 previous dose was PRP-T (ActHib®, Pentacel®, Hiberix®), Vaxelis® or unknown <b>8 weeks and age 12 through 59 months (as final dose)</b> if current age is younger than 12 months <b>and</b> first dose was administered at age 7 through 11 months; OR if current age is 12 through 59 months <b>and</b> first dose was administered before the 1 <sup>st</sup> birthday <b>and</b> second dose was administered at younger than 15 months; OR if both doses were PedvaxiHIB® and were administered before the 1st birthday	<b>8 weeks (as final dose)</b> This dose only necessary for children age 12 through 59 months who received 3 doses before the 1 <sup>st</sup> birthday.	
Pneumococcal conjugate	6 weeks	<b>No further doses needed</b> for healthy children if first dose was administered at age 24 months or older. <b>4 weeks</b> if first dose was administered before the 1 <sup>st</sup> birthday <b>8 weeks (as final dose for healthy children)</b> if first dose was administered at the 1 <sup>st</sup> birthday or after	<b>No further doses needed</b> for healthy children if previous dose was administered at age 24 months or older <b>4 weeks</b> if current age is younger than 12 months and previous dose was administered at <7 months old <b>8 weeks (as final dose for healthy children)</b> if previous dose was administered between 7–11 months (wait until at least 12 months old); OR if current age is 12 months or older and at least 1 dose was administered before age 12 months	<b>8 weeks (as final dose)</b> This dose only necessary for children age 12 through 59 months who received 3 doses before age 12 months or for children at high risk who received 3 doses at any age.	
Inactivated poliovirus	6 weeks	<b>4 weeks</b>	<b>4 weeks</b> if current age is <4 years <b>6 months (as final dose)</b> if current age is 4 years or older	<b>6 months (minimum age 4 years for final dose)</b>	
Measles, mumps, rubella	12 months	<b>4 weeks</b>			
Varicella	12 months	<b>3 months</b>			
Hepatitis A	12 months	<b>6 months</b>			
Meningococcal ACWY	2 months MenACWY-CRM 9 months MenACWY-D 2 years MenACWY-TT	<b>8 weeks</b>	See Notes	See Notes	
Children and adolescents age 7 through 18 years					
Meningococcal ACWY	Not applicable (N/A)	<b>8 weeks</b>			
Tetanus, diphtheria, and acellular pertussis	7 years	<b>4 weeks</b>	<b>4 weeks</b> if first dose of DTaP/DT was administered before the 1 <sup>st</sup> birthday <b>6 months (as final dose)</b> if first dose of DTaP/DT or Tdap/Td was administered at or after the 1 <sup>st</sup> birthday	<b>6 months</b> if first dose of DTaP/DT was administered before the 1 <sup>st</sup> birthday	
Human papillomavirus	9 years	<b>Routine dosing intervals are recommended.</b>			
Hepatitis A	N/A	<b>6 months</b>			
Hepatitis B	N/A	<b>4 weeks</b>	<b>8 weeks and at least 16 weeks after first dose</b>		
Inactivated poliovirus	N/A	<b>4 weeks</b>	<b>6 months</b> A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose.	A fourth dose of IPV is indicated if all previous doses were administered at <4 years or if the third dose was administered <6 months after the second dose.	
Measles, mumps, rubella	N/A	<b>4 weeks</b>			
Varicella	N/A	<b>3 months</b> if younger than age 13 years. <b>4 weeks</b> if age 13 years or older			
Dengue	9 years	<b>6 months</b>	6 months		

**Table 3**

## Recommended Child and Adolescent Immunization Schedule by Medical Indication, United States, 2022

Always use this table in conjunction with Table 1 and the Notes that follow.

VACCINE	INDICATION									
	Pregnancy	Immunocompromised status (excluding HIV infection)	HIV infection CD4+ count <sup>1</sup>		Kidney failure, end-stage renal disease, or on hemodialysis	Heart disease or chronic lung disease	CSF leak or cochlear implant	Asplenia or persistent complement component deficiencies	Chronic liver disease	Diabetes
			<15% or total CD4 cell count of <200/mm <sup>3</sup>	≥15% and total CD4 cell count of ≥200/mm <sup>3</sup>						
Hepatitis B										
Rotavirus		SCID <sup>2</sup>								
Diphtheria, tetanus, and acellular pertussis (DTaP)										
<i>Haemophilus influenzae</i> type b										
Pneumococcal conjugate										
Inactivated poliovirus										
Influenza (IV4) or Influenza (LAIV4)										
Influenza (LAIV4)						Asthma, wheezing: 2–4yrs <sup>3</sup>				
Measles, mumps, rubella	*									
Varicella	*									
Hepatitis A										
Tetanus, diphtheria, and acellular pertussis (Tdap)										
Human papillomavirus	*									
Meningococcal ACWY										
Meningococcal B										
Pneumococcal polysaccharide										
Dengue										

Vaccination according to the routine schedule recommended
Recommended for persons with an additional risk factor for which the vaccine would be indicated
Vaccination is recommended, and additional doses may be necessary based on medical condition or vaccine. See Notes.
Precaution—vaccine might be indicated if benefit of protection outweighs risk of adverse reaction
Contraindicated or not recommended—vaccine should not be administered  
\*Vaccinate after pregnancy
No recommendation/not applicable

1 For additional information regarding HIV laboratory parameters and use of live vaccines, see the *General Best Practice Guidelines for Immunization*, "Altered Immunocompetence," at [www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html](http://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html) and Table 4-1 (footnote J) at [www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html](http://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html).  
<sup>2</sup> Severe Combined Immunodeficiency

## Child and Adolescents Immunization

Revision #5

Created 12 February 2022 21:20:49 by PHILLIP KIM

Updated 4 April 2022 05:03:55 by Katarina Soewono