

Tachycardia

Narrow complex tachycardia

Evaluation

- Obtain EKG and complete set of vital sign to check for hemodynamic instability
- If unstable follow ACLS protocol

Management

- Unstable: follow ACLS protocol and do not delay synchronized cardioversion
- Try vagal maneuver or adenosine.
- Vagal maneuver:
 - Valsalva
 - Carotid massage
- Adenosine: Warn patients that adenosine will make them feel terrible but it will be short.
 - Dosing:
 - Peripheral IV 6mg x1. may repeat one more time with the same dose several minutes later if not improving, then increase to 12mg dose x1
 - Central line: 3mg with a repeat dose and then increased dose to 6mg
 - **DO NOT use in heart transplant recipient, wide complex tachycardia and WPW or previous allergy**

Wide Complex Tachycardia

Evaluation

- Is patient stable or unstable
- If unstable or in doubt call code blue and follow ACLS protocol
- If stable obtain EKG, troponin, and electrolyte panel including magnesium
- Look for precipitating cause and medication list for QT prolonging agents

Management

- Replete electrolytes

- Discontinue QT prolonged agents. Magnesium sulfate 2g IV should be given if patient have prolonged Qtc (>450 in women and >470 in men)

Revision #2

Created 25 February 2022 06:11:29 by Katarina Soewono

Updated 13 June 2022 07:01:34 by Katarina Soewono