

Twins

At risk for hyperemesis, GDM, HTN, hemorrhage, cesarean, postpartum depression, preeclampsia, preterm delivery, IUGR, birth/genetic defects

- Pre-term labor education
- Fetal kick count starting at 28 weeks (10 kicks/movement within 2hrs or less)
- NST weekly starting at 36 weeks for uncomplicated dichorionic
- NST weekly starting at 32 weeks for uncomplicated monochorionic diamniotic.

Management

- Dichorionic twins
 - Early US for chorionicity, detailed US at 18-22 wks, growth every 4-6 weeks, delivery at 38 weeks
- Monochorionic diamniotic twin
 - Early US for chorionicity, US for TTTS (twin to twin transfusion syndrome) f/u every 2 weeks at 16 weeks, growth every 4 weeks, delivery at 34-37 weeks
- Consults with perinatologist for discordant growth, monoamniotic twins (cord accidents) or monochorionic twins (twin to twin transfusions)

Pocket Obstetrics and gynecology by K Joseph Hurt 2nd edition

<https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2021/06/indications-for-outpatient-antenatal-fetal-surveillance>

Revision #4

Created 25 February 2022 06:51:05 by Katarina Soewono

Updated 30 May 2022 05:19:48 by Katarina Soewono